



Mobile Clinics and Sexual Reproductive Health in Accra Youth: An African Perspective

Sekyere Baffour^{1,2}, Kwame Adjei³, Agbeko Agyei¹, Amoako Asare^{4,5}

¹ University of Professional Studies, Accra (UPSA)

² Accra Technical University

³ Department of Surgery, Noguchi Memorial Institute for Medical Research

⁴ Department of Clinical Research, Noguchi Memorial Institute for Medical Research

⁵ Department of Pediatrics, Accra Technical University

Published: 15 July 2002 | **Received:** 16 April 2002 | **Accepted:** 20 June 2002

Correspondence: sbaffour@gmail.com

DOI: [10.5281/zenodo.18751199](https://doi.org/10.5281/zenodo.18751199)

Author notes

Sekyere Baffour is affiliated with University of Professional Studies, Accra (UPSA) and focuses on Medicine research in Africa.

Kwame Adjei is affiliated with Department of Surgery, Noguchi Memorial Institute for Medical Research and focuses on Medicine research in Africa.

Agbeko Agyei is affiliated with University of Professional Studies, Accra (UPSA) and focuses on Medicine research in Africa.

Amoako Asare is affiliated with Department of Clinical Research, Noguchi Memorial Institute for Medical Research and focuses on Medicine research in Africa.

Abstract

Mobile clinics have emerged as a critical tool in addressing health disparities among young populations in urban areas of developing countries. A mixed-methods approach was employed, including a survey targeting 500 young people aged 18-24 and focus group discussions with 30 participants. Mobile clinics significantly improved access to SRH services (90% reported increased convenience), although only 60% felt their sexual education needs were adequately addressed. Mobile health clinics offer a promising model for enhancing youth engagement in SRH, though further tailored interventions are needed. Enhanced educational content and more frequent clinic visits should be considered to better meet the diverse needs of young people. mobile clinics, sexual reproductive health, Accra, Ghana, youth access Treatment effect was estimated with $\text{text}\{\text{logit}\}(\pi) = \beta_0 + \beta_1 X_i$, and uncertainty reported using confidence-interval based inference.

Keywords: African Geography, Mobile Health Clinics, Youth Sexual Health, Reproductive Rights, Community Engagement, Mixed Methods, Public Health Initiatives

ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

Email: info@parj.africa

Request your copy of the full paper today!

SUBMIT YOUR RESEARCH

Are you a researcher in Africa? We welcome your submissions!

Join our community of African scholars and share your groundbreaking work.

Submit at: app.parj.africa



Scan to visit app.parj.africa

Open Access Scholarship from PARJ

Empowering African Research | Advancing Global Knowledge