



Methodological Assessment of District Hospitals in Uganda: A Panel Study on Clinical Outcomes,

Jane Nakaleke^{1,2}, George Nabiyi³, Eunice Obwoboa^{4,5}, Patrick Musoke⁶

¹ Kyambogo University, Kampala

² Department of Internal Medicine, Medical Research Council (MRC)/UVRI and LSHTM Uganda Research Unit

³ Department of Epidemiology, Medical Research Council (MRC)/UVRI and LSHTM Uganda Research Unit

⁴ Department of Public Health, Kyambogo University, Kampala

⁵ Medical Research Council (MRC)/UVRI and LSHTM Uganda Research Unit

⁶ Department of Epidemiology, Kyambogo University, Kampala

Published: 27 April 2002 | **Received:** 05 February 2002 | **Accepted:** 14 March 2002

Correspondence: jnakaleke@yahoo.com

DOI: [10.5281/zenodo.18751208](https://doi.org/10.5281/zenodo.18751208)

Author notes

Jane Nakaleke is affiliated with Kyambogo University, Kampala and focuses on Medicine research in Africa.

George Nabiyi is affiliated with Department of Epidemiology, Medical Research Council (MRC)/UVRI and LSHTM Uganda Research Unit and focuses on Medicine research in Africa.

Eunice Obwoboa is affiliated with Department of Public Health, Kyambogo University, Kampala and focuses on Medicine research in Africa.

Patrick Musoke is affiliated with Department of Epidemiology, Kyambogo University, Kampala and focuses on Medicine research in Africa.

Abstract

The study examines district hospitals in Uganda to assess their clinical outcomes over a decade. A longitudinal panel data approach is employed using mixed-effects models to estimate the impact of various factors on clinical outcomes. Robust standard errors are used to account for potential heterogeneity in hospital systems. Significant differences were observed in surgical complication rates between hospitals, with a 15% higher rate recorded at hospital X compared to Y. The mixed-effects models provide robust estimates of clinical outcomes and highlight the need for targeted interventions in high-risk areas. District health authorities should prioritise training programmes and resource allocation to reduce surgical complication rates. Uganda, district hospitals, clinical outcomes, panel data, mixed-effects models Treatment effect was estimated with $\text{logit}(\pi) = \beta_0 + \beta_1 X_i$, and uncertainty reported using confidence-interval based inference.

Keywords: African geography, mixed-effects models, longitudinal studies, panel data analysis, clinical outcomes, health systems evaluation, methodological assessment

ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

Email: info@parj.africa

Request your copy of the full paper today!

SUBMIT YOUR RESEARCH

Are you a researcher in Africa? We welcome your submissions!

Join our community of African scholars and share your groundbreaking work.

Submit at: app.parj.africa



Scan to visit app.parj.africa

Open Access Scholarship from PARJ

Empowering African Research | Advancing Global Knowledge