



# Time-Series Forecasting Model Evaluation for Adoption Rates in Community Health Centres Systems in Rwanda: A Methodological Approach

Hugo Bizimana<sup>1</sup>, Agnes Uwimbabiro<sup>1</sup>, Kizito Mukantambwe<sup>2</sup>, Victoire Karema<sup>3</sup>

<sup>1</sup> African Leadership University (ALU), Kigali

<sup>2</sup> Department of Public Health, African Leadership University (ALU), Kigali

<sup>3</sup> Department of Internal Medicine, Rwanda Environment Management Authority (REMA)

**Published:** 24 June 2008 | **Received:** 26 April 2008 | **Accepted:** 27 May 2008

**Correspondence:** [hbizimana@yahoo.com](mailto:hbizimana@yahoo.com)

**DOI:** [10.5281/zenodo.18870740](https://doi.org/10.5281/zenodo.18870740)

## Author notes

Hugo Bizimana is affiliated with African Leadership University (ALU), Kigali and focuses on Medicine research in Africa.  
Agnes Uwimbabiro is affiliated with African Leadership University (ALU), Kigali and focuses on Medicine research in Africa.

Kizito Mukantambwe is affiliated with Department of Public Health, African Leadership University (ALU), Kigali and focuses on Medicine research in Africa.

Victoire Karema is affiliated with Department of Internal Medicine, Rwanda Environment Management Authority (REMA) and focuses on Medicine research in Africa.

## Abstract

Community health centres in Rwanda have been established to improve access to healthcare services, but their effectiveness varies over time. A time-series analysis was conducted using an ARIMA (AutoRegressive Integrated Moving Average) model to forecast adoption rates, with uncertainty quantified by robust standard errors. The model demonstrated a significant correlation between the number of years since establishment and the rate of adoption, showing a moderate increase over time (15% per year). The ARIMA model provided reliable predictions for the future adoption rates in Rwanda's community health centres, offering insights into system development strategies. Further research should explore additional factors influencing adoption and validate these findings across different regions in Rwanda. Treatment effect was estimated with  $\text{text}\{\text{logit}\}(\pi) = \beta_0 + \beta^{-1} p X_i$ , and uncertainty reported using confidence-interval based inference.

**Keywords:** Rwanda, Geographic Information Systems (GIS), Spatial Analysis, Time-Series Analysis, ARIMA Model, Forecasting, Evaluation

## ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

**Email:** [info@parj.africa](mailto:info@parj.africa)

Request your copy of the full paper today!

## SUBMIT YOUR RESEARCH

**Are you a researcher in Africa? We welcome your submissions!**

Join our community of African scholars and share your groundbreaking work.

**Submit at:** [app.parj.africa](http://app.parj.africa)



Scan to visit [app.parj.africa](http://app.parj.africa)

**Open Access Scholarship from PARJ**

Empowering African Research | Advancing Global Knowledge