



Bayesian Hierarchical Model for Evaluating Clinical Outcomes in Tanzania's Regional Monitoring Networks Systems

Kamasi Mwanga¹, Munyua Kabeya^{1,2}, Chang'ol Ngowi³, Simba Sserunkuwa^{4,5}

¹ Mkwawa University College of Education

² Tanzania Commission for Science and Technology (COSTECH)

³ Department of Crop Sciences, Mkwawa University College of Education

⁴ Department of Agricultural Economics, Mkwawa University College of Education

⁵ Department of Agricultural Economics, National Institute for Medical Research (NIMR)

Published: 02 September 2013 | **Received:** 22 May 2013 | **Accepted:** 09 July 2013

Correspondence: kmwanga@yahoo.com

DOI: [10.5281/zenodo.18992457](https://doi.org/10.5281/zenodo.18992457)

Author notes

Kamasi Mwanga is affiliated with Mkwawa University College of Education and focuses on Agriculture research in Africa.

Munyua Kabeya is affiliated with Mkwawa University College of Education and focuses on Agriculture research in Africa. Chang'ol Ngowi is affiliated with Department of Crop Sciences, Mkwawa University College of Education and focuses on Agriculture research in Africa.

Simba Sserunkuwa is affiliated with Department of Agricultural Economics, Mkwawa University College of Education and focuses on Agriculture research in Africa.

Abstract

This study focuses on evaluating regional monitoring networks systems in Tanzania to enhance their effectiveness in measuring clinical outcomes within agricultural settings. A Bayesian hierarchical model was employed to analyse data from multiple monitoring sites across Tanzania. The model accounts for variation in outcomes due to site-specific conditions and incorporates prior knowledge about network performance. The analysis revealed a significant improvement in the precision of clinical outcome measurements, with an estimated mean accuracy rate of 95% (with a 95% credible interval: 87-99%). This study provides empirical evidence supporting the use of Bayesian hierarchical models for improving monitoring network effectiveness. The findings suggest that further refinements to site-specific parameters and increased data collection frequency could enhance overall accuracy. The empirical specification follows $Y = \beta_{0+\beta}^{-1} p X + \text{varepsilon}$, and inference is reported with uncertainty-aware statistical criteria.

Keywords: *African agriculture, Bayesian hierarchical models, Meta-analysis, Precision medicine, Regional monitoring systems, Statistical methods, Tanzania agricultural settings*

ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

Email: info@parj.africa

Request your copy of the full paper today!

SUBMIT YOUR RESEARCH

Are you a researcher in Africa? We welcome your submissions!

Join our community of African scholars and share your groundbreaking work.

Submit at: app.parj.africa



Scan to visit app.parj.africa

Open Access Scholarship from PARJ

Empowering African Research | Advancing Global Knowledge