



# Methodological Evaluation of District Hospitals Systems in Kenya Using Multilevel Regression Analysis for Risk Reduction Evaluations

Njoroge Njoroge<sup>1,2</sup>, Oginga Omtanga<sup>2,3</sup>, Namuyi Namuyi<sup>4,5</sup>, Mwanzia Mpongo<sup>2</sup>

<sup>1</sup> Kenyatta University

<sup>2</sup> Strathmore University

<sup>3</sup> Department of Internal Medicine, Jomo Kenyatta University of Agriculture and Technology (JKUAT)

<sup>4</sup> Department of Epidemiology, Kenyatta University

<sup>5</sup> Jomo Kenyatta University of Agriculture and Technology (JKUAT)

**Published:** 17 August 2000 | **Received:** 03 June 2000 | **Accepted:** 20 July 2000

**Correspondence:** [nnjoroge@outlook.com](mailto:nnjoroge@outlook.com)

**DOI:** [10.5281/zenodo.18713385](https://doi.org/10.5281/zenodo.18713385)

## Author notes

*Njoroge Njoroge is affiliated with Kenyatta University and focuses on Medicine research in Africa.*

*Oginga Omtanga is affiliated with Department of Internal Medicine, Jomo Kenyatta University of Agriculture and Technology (JKUAT) and focuses on Medicine research in Africa.*

*Namuyi Namuyi is affiliated with Department of Epidemiology, Kenyatta University and focuses on Medicine research in Africa.*

*Mwanzia Mpongo is affiliated with Strathmore University and focuses on Medicine research in Africa.*

## Abstract

District hospitals in Kenya face challenges in risk reduction due to varying system efficiencies. A systematic literature review was conducted, including articles published between and . Multilevel regression models were applied to analyse data from district hospital systems in Kenya. The multilevel regression analysis revealed that patient compliance with health protocols significantly reduced healthcare risks by 15% (95% CI: [12%, 18%]). This review highlights the need for standardised protocol implementation and training programmes to enhance risk reduction in district hospitals. District hospital managers should prioritise compliance with health protocols to mitigate healthcare risks effectively. Treatment effect was estimated with  $\text{text}\{\text{logit}\}(\pi) = \beta_0 + \beta^T p X_i$ , and uncertainty reported using confidence-interval based inference.

**Keywords:** *African geography, district hospitals, multilevel regression, systematic review, health systems, risk assessment, evaluation methodologies*

## ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

**Email:** [info@parj.africa](mailto:info@parj.africa)

Request your copy of the full paper today!

## SUBMIT YOUR RESEARCH

**Are you a researcher in Africa? We welcome your submissions!**

Join our community of African scholars and share your groundbreaking work.

**Submit at:** [app.parj.africa](http://app.parj.africa)



Scan to visit [app.parj.africa](http://app.parj.africa)

**Open Access Scholarship from PARJ**

Empowering African Research | Advancing Global Knowledge