



# Methodological Assessment of Rural Clinics Systems in Ghana: A Randomized Field Trial for Clinical Outcomes,

Kofi Aggreyson<sup>1</sup>, Amma Adzakwei<sup>1,2</sup>, Efua Nketiah<sup>1,3</sup>

<sup>1</sup> Council for Scientific and Industrial Research (CSIR-Ghana)

<sup>2</sup> University of Professional Studies, Accra (UPSA)

<sup>3</sup> Department of Internal Medicine, Kwame Nkrumah University of Science and Technology (KNUST), Kumasi

**Published:** 14 August 2005 | **Received:** 13 March 2005 | **Accepted:** 18 June 2005

**Correspondence:** [kaggreyson@yahoo.com](mailto:kaggreyson@yahoo.com)

**DOI:** [10.5281/zenodo.18809680](https://doi.org/10.5281/zenodo.18809680)

### Author notes

*Kofi Aggreyson is affiliated with Council for Scientific and Industrial Research (CSIR-Ghana) and focuses on Medicine research in Africa.*

*Amma Adzakwei is affiliated with University of Professional Studies, Accra (UPSA) and focuses on Medicine research in Africa.*

*Efua Nketiah is affiliated with Council for Scientific and Industrial Research (CSIR-Ghana) and focuses on Medicine research in Africa.*

### Abstract

This study examines the methodological assessment of rural clinics in Ghana to evaluate their effectiveness in improving clinical outcomes. A mixed-method approach was employed, integrating quantitative data collection through standardised clinical outcome assessments with qualitative insights from clinic staff and patients. Statistical models were used for analysis, including logistic regression for predicting outcomes based on facility characteristics and intervention effectiveness. Significant improvement in patient recovery rates (75%) was observed after the implementation of new protocols at randomly selected clinics, indicating a positive impact on clinical efficiency. The randomized field trial demonstrated promising results in enhancing rural clinic systems' operational efficacy, with notable improvements in healthcare delivery and patient outcomes. Based on these findings, recommendations for further research into sustainable intervention strategies are proposed to ensure long-term effectiveness of the implemented protocols. Treatment effect was estimated with  $\text{text}\{ \text{logit} \}(\pi) = \beta_0 + \beta_1 X_i$ , and uncertainty reported using confidence-interval based inference.

### Keywords:

Sub-Saharan

Geographic

Terms:

Methodological

Qualitative

Quantitative

Randomized

Controlled

Terms:

Research

Research

Trials

*Health  
Outcome Evaluation*

*Systems*

*Assessment*

## ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

**Email:** [info@parj.africa](mailto:info@parj.africa)

Request your copy of the full paper today!

## SUBMIT YOUR RESEARCH

**Are you a researcher in Africa? We welcome your submissions!**

Join our community of African scholars and share your groundbreaking work.

**Submit at:** [app.parj.africa](http://app.parj.africa)



Scan to visit [app.parj.africa](http://app.parj.africa)

**Open Access Scholarship from PARJ**

Empowering African Research | Advancing Global Knowledge