



# Methodological Assessment of District Hospitals Systems in Nigeria: Quasi-Experimental Design for Risk Reduction Evaluation

Osita Ifidon<sup>1,2</sup>, Nkechi Emenike<sup>3</sup>, Chinwe Okpara<sup>1,4</sup>, Sunday Ofoh<sup>2,5</sup>

<sup>1</sup> Nigerian Institute of Social and Economic Research (NISER)

<sup>2</sup> Obafemi Awolowo University, Ile-Ife

<sup>3</sup> Department of Clinical Research, Obafemi Awolowo University, Ile-Ife

<sup>4</sup> Usmanu Danfodiyo University, Sokoto

<sup>5</sup> Department of Internal Medicine, Usmanu Danfodiyo University, Sokoto

**Published:** 16 February 2009 | **Received:** 19 September 2008 | **Accepted:** 30 December 2008

**Correspondence:** [oifidon@gmail.com](mailto:oifidon@gmail.com)

**DOI:** [10.5281/zenodo.18886894](https://doi.org/10.5281/zenodo.18886894)

## Author notes

*Osita Ifidon is affiliated with Nigerian Institute of Social and Economic Research (NISER) and focuses on Medicine research in Africa.*

*Nkechi Emenike is affiliated with Department of Clinical Research, Obafemi Awolowo University, Ile-Ife and focuses on Medicine research in Africa.*

*Chinwe Okpara is affiliated with Usmanu Danfodiyo University, Sokoto and focuses on Medicine research in Africa.  
Sunday Ofoh is affiliated with Department of Internal Medicine, Usmanu Danfodiyo University, Sokoto and focuses on Medicine research in Africa.*

## Abstract

Nigeria's district hospitals face significant challenges in managing their healthcare systems efficiently and effectively. A mixed-methods approach combining quantitative data analysis with qualitative interviews to evaluate the efficiency and effectiveness of healthcare delivery systems in selected districts. District hospitals reported an average reduction in patient wait times by 25% after implementing new scheduling protocols, though variability existed among different regions. The quasi-experimental design revealed promising initial outcomes for risk reduction but also highlighted ongoing challenges that require further investigation and intervention strategies. Further research should include a longitudinal study to monitor the sustainability of implemented changes and explore additional cost-effective solutions. Treatment effect was estimated with  $\text{logit}(\pi) = \beta_0 + \beta_1 X_i$ , and uncertainty reported using confidence-interval based inference.

**Keywords:** *Sub-Saharan, health policy, mixed-methods, quasi-experimental, evaluation, decentralization, intervention effectiveness*

## ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

**Email:** [info@parj.africa](mailto:info@parj.africa)

Request your copy of the full paper today!

## SUBMIT YOUR RESEARCH

**Are you a researcher in Africa? We welcome your submissions!**

Join our community of African scholars and share your groundbreaking work.

**Submit at:** [app.parj.africa](http://app.parj.africa)



Scan to visit [app.parj.africa](http://app.parj.africa)

**Open Access Scholarship from PARJ**

Empowering African Research | Advancing Global Knowledge