



Methodological Assessment of District Hospital Systems in South Africa: Panel Data Estimation for Adoption Rates Analysis

Makana Khumalo¹, Siphon Mabotsa², Nokuthula Ngwenya^{3,4}

¹ Department of Pediatrics, North-West University

² University of Zululand

³ Department of Surgery, University of Zululand

⁴ Tshwane University of Technology (TUT)

Published: 12 June 2007 | **Received:** 16 March 2007 | **Accepted:** 12 May 2007

Correspondence: mkhumalo@outlook.com

DOI: [10.5281/zenodo.18845089](https://doi.org/10.5281/zenodo.18845089)

Author notes

Makana Khumalo is affiliated with Department of Pediatrics, North-West University and focuses on Medicine research in Africa.

Siphon Mabotsa is affiliated with University of Zululand and focuses on Medicine research in Africa.

Nokuthula Ngwenya is affiliated with Department of Surgery, University of Zululand and focuses on Medicine research in Africa.

Abstract

District hospitals in South Africa play a crucial role in providing healthcare services to underserved populations. However, their operational efficiency and resource allocation vary significantly across different districts. A mixed-method approach was employed, integrating quantitative panel-data econometric techniques with qualitative interviews. The dataset includes longitudinal patient records and administrative health information system (HIS) data from ten districts over a five-year period. The analysis revealed that the adoption rate for antiretroviral therapy (ART) in district hospitals varied significantly, ranging from 70% to 95%, with some districts showing lower adherence rates due to resource limitations and patient education gaps. This study underscores the importance of tailored interventions to improve ART adoption rates and highlights the need for continuous monitoring and support mechanisms within district hospital systems. Healthcare policymakers should prioritise funding, staff training, and community engagement strategies in underserved districts to enhance healthcare service delivery and patient outcomes. district hospitals, South Africa, panel data analysis, adoption rates, antiretroviral therapy Treatment effect was estimated with $\text{text}\{\text{logit}\}(\pi) = \beta_0 + \beta_1 X_i$, and uncertainty reported using confidence-interval based inference.

Keywords: Sub-Saharan, African, Spatial, Econometrics, Healthcare, Systems, Geography, Panel

ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

Email: info@parj.africa

Request your copy of the full paper today!

SUBMIT YOUR RESEARCH

Are you a researcher in Africa? We welcome your submissions!

Join our community of African scholars and share your groundbreaking work.

Submit at: app.parj.africa



Scan to visit app.parj.africa

Open Access Scholarship from PARJ

Empowering African Research | Advancing Global Knowledge