



# Methodological Evaluation of District Hospitals' Clinical Outcomes in Kenya Using Difference-in-Differences Analysis

Karurvy Kinyanjui<sup>1,2</sup>, Kamau Otieno<sup>1,2</sup>, Oyunga Ngugi<sup>3,4</sup>, Rutagwira Ochieng<sup>5,6</sup>

<sup>1</sup> Pwani University

<sup>2</sup> Strathmore University

<sup>3</sup> Egerton University

<sup>4</sup> Maseno University

<sup>5</sup> Department of Epidemiology, Strathmore University

<sup>6</sup> Department of Internal Medicine, Egerton University

Published: 02 January 2002 | Received: 16 August 2001 | Accepted: 22 November 2001

Correspondence: [kkinyanjui@gmail.com](mailto:kkinyanjui@gmail.com)

DOI: [10.5281/zenodo.18739624](https://doi.org/10.5281/zenodo.18739624)

### Author notes

*Karurvy Kinyanjui is affiliated with Pwani University and focuses on Medicine research in Africa.*

*Kamau Otieno is affiliated with Strathmore University and focuses on Medicine research in Africa.*

*Oyunga Ngugi is affiliated with Egerton University and focuses on Medicine research in Africa.*

*Rutagwira Ochieng is affiliated with Department of Epidemiology, Strathmore University and focuses on Medicine research in Africa.*

### Abstract

District hospitals in Kenya play a critical role in healthcare delivery, particularly for underserved rural populations. However, their clinical outcomes are often suboptimal due to various systemic issues. Clinical outcome data from a sample of district hospitals in Kenya were analysed over two time periods: before and after implementing quality improvement measures. DiD analysis was employed to estimate treatment effects, accounting for unobserved heterogeneity. The difference-in-differences model revealed a statistically significant positive impact on patient recovery rates ( $p < 0.05$ ) in the intervention group compared to controls. Our analysis supports the efficacy of DiD modelling as a robust method for evaluating healthcare system improvements, with notable benefits observed in district hospital clinical outcomes. Future studies should consider expanding the DiD model application and incorporating additional variables to enhance its predictive accuracy. Treatment effect was estimated with  $\text{text}\{logit\}(\pi) = \beta_0 + \beta_1 X_i$ , and uncertainty reported using confidence-interval based inference.

**Keywords:** Kenya, District Hospitals, Clinical Outcomes, Difference-in-Differences, Evaluation, Metrics, Analytics

## ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

**Email:** [info@parj.africa](mailto:info@parj.africa)

Request your copy of the full paper today!

## SUBMIT YOUR RESEARCH

**Are you a researcher in Africa? We welcome your submissions!**

Join our community of African scholars and share your groundbreaking work.

**Submit at:** [app.parj.africa](http://app.parj.africa)



Scan to visit [app.parj.africa](http://app.parj.africa)

**Open Access Scholarship from PARJ**

Empowering African Research | Advancing Global Knowledge