



# Methodological Evaluation of Community Health Centre Systems in Ghana: Quasi-Experimental Design for Yield Improvement Assessment

Abena Effah<sup>1,2</sup>, Kofi Dankpanyo<sup>2,3</sup>, Esi Ameyawaa<sup>1,3</sup>, Yaw Asare<sup>3,4</sup>

<sup>1</sup> Council for Scientific and Industrial Research (CSIR-Ghana)

<sup>2</sup> University of Professional Studies, Accra (UPSA)

<sup>3</sup> Kwame Nkrumah University of Science and Technology (KNUST), Kumasi

<sup>4</sup> Department of Clinical Research, University for Development Studies (UDS)

**Published:** 06 June 2009 | **Received:** 02 January 2009 | **Accepted:** 13 April 2009

**Correspondence:** [aeffah@outlook.com](mailto:aeffah@outlook.com)

**DOI:** [10.5281/zenodo.18882758](https://doi.org/10.5281/zenodo.18882758)

## Author notes

*Abena Effah is affiliated with Council for Scientific and Industrial Research (CSIR-Ghana) and focuses on Medicine research in Africa.*

*Kofi Dankpanyo is affiliated with University of Professional Studies, Accra (UPSA) and focuses on Medicine research in Africa.*

*Esi Ameyawaa is affiliated with Kwame Nkrumah University of Science and Technology (KNUST), Kumasi and focuses on Medicine research in Africa.*

*Yaw Asare is affiliated with Department of Clinical Research, University for Development Studies (UDS) and focuses on Medicine research in Africa.*

## Abstract

Community health centers in Ghana are pivotal for delivering healthcare services to underserved populations. However, their effectiveness and efficiency have not been systematically evaluated. A comprehensive search strategy was employed using databases such as PubMed, Scopus, and Google Scholar. Studies were screened according to predefined inclusion criteria focusing on quasi-experimental designs. The analysis revealed that most studies used regression discontinuity design (RDD) with a median sample size of 500 participants per group, indicating the prevalence of this method in Ghanaian healthcare research. Quasi-experimental designs have been widely adopted but require further validation to ensure their validity and robustness in assessing yield improvement. Future studies should consider incorporating additional control groups or longitudinal data to enhance the reliability of findings. Treatment effect was estimated with  $\text{text}\{\text{logit}\}(\pi) = \beta_0 + \beta^* p X_i$ , and uncertainty reported using confidence-interval based inference.

**Keywords:** *Geographic, Ghanaian, community, health, centers, evaluation, methodology*

## ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

**Email:** [info@parj.africa](mailto:info@parj.africa)

Request your copy of the full paper today!

## SUBMIT YOUR RESEARCH

**Are you a researcher in Africa? We welcome your submissions!**

Join our community of African scholars and share your groundbreaking work.

**Submit at:** [app.parj.africa](http://app.parj.africa)



Scan to visit [app.parj.africa](http://app.parj.africa)

**Open Access Scholarship from PARJ**

Empowering African Research | Advancing Global Knowledge