



# Methodological Evaluation of Rural Clinics in Tanzania Using Time-Series Forecasting Models for Clinical Outcome Measurement,

Kamali Mwakwaya<sup>1</sup>, Mkono Simiyu<sup>2,3</sup>

<sup>1</sup> Nelson Mandela African Institution of Science and Technology (NM-AIST), Arusha

<sup>2</sup> University of Dar es Salaam

<sup>3</sup> Department of Public Health, Nelson Mandela African Institution of Science and Technology (NM-AIST), Arusha

**Published:** 28 July 2005 | **Received:** 14 March 2005 | **Accepted:** 04 July 2005

**Correspondence:** [kmwakwaya@aol.com](mailto:kmwakwaya@aol.com)

**DOI:** [10.5281/zenodo.18806523](https://doi.org/10.5281/zenodo.18806523)

## Author notes

*Kamali Mwakwaya is affiliated with Nelson Mandela African Institution of Science and Technology (NM-AIST), Arusha and focuses on Medicine research in Africa.*

*Mkono Simiyu is affiliated with University of Dar es Salaam and focuses on Medicine research in Africa.*

## Abstract

This study addresses a current research gap in Medicine concerning Methodological evaluation of rural clinics systems in Tanzania: time-series forecasting model for measuring clinical outcomes in Tanzania. The objective is to formulate a rigorous model, state verifiable assumptions, and derive results with direct analytical or practical implications. A mixed-methods design was used, combining survey and interview data collected over the study period. The results establish bounded error under perturbation, a convergent estimation process under stated assumptions, and a stable link between the proposed metric and observed outcomes. The findings provide a reproducible analytical basis for subsequent theoretical and applied extensions. Stakeholders should prioritise inclusive, locally grounded strategies and improve data transparency. Methodological evaluation of rural clinics systems in Tanzania: time-series forecasting model for measuring clinical outcomes, Tanzania, Africa, Medicine, longitudinal study This work contributes a formal specification, transparent assumptions, and mathematically interpretable claims. Treatment effect was estimated with  $\text{text}\{logit\}(\pi) = \beta_0 + \beta^T p X_i$ , and uncertainty reported using confidence-interval based inference.

**Keywords:** *Rural health, Tanzania, time-series analysis, forecasting models, clinical outcomes, longitudinal studies, health systems assessment*

## ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

**Email:** [info@parj.africa](mailto:info@parj.africa)

Request your copy of the full paper today!

## SUBMIT YOUR RESEARCH

**Are you a researcher in Africa? We welcome your submissions!**

Join our community of African scholars and share your groundbreaking work.

**Submit at:** [app.parj.africa](http://app.parj.africa)



Scan to visit [app.parj.africa](http://app.parj.africa)

**Open Access Scholarship from PARJ**

Empowering African Research | Advancing Global Knowledge