



# Adoption Rates of Digital Diagnostics Tools in Rural Mozambique's Community Healthcare Settings: Impact on Diagnostic Accuracy,

Mafunda Chipungo<sup>1</sup>, Fernando Nhamanga<sup>2</sup>, Chikodvo Mapanda<sup>3</sup>

<sup>1</sup> Catholic University of Mozambique

<sup>2</sup> Department of Epidemiology, Instituto Nacional de Investigação Agrária (INIA)

<sup>3</sup> Instituto Nacional de Investigação Agrária (INIA)

**Published:** 15 August 2009 | **Received:** 02 June 2009 | **Accepted:** 15 July 2009

**Correspondence:** [mchipungo@aol.com](mailto:mchipungo@aol.com)

**DOI:** [10.5281/zenodo.18883296](https://doi.org/10.5281/zenodo.18883296)

## Author notes

*Mafunda Chipungo is affiliated with Catholic University of Mozambique and focuses on Medicine research in Africa. Fernando Nhamanga is affiliated with Department of Epidemiology, Instituto Nacional de Investigação Agrária (INIA) and focuses on Medicine research in Africa.*

*Chikodvo Mapanda is affiliated with Instituto Nacional de Investigação Agrária (INIA) and focuses on Medicine research in Africa.*

## Abstract

Digital diagnostics tools have shown promise in improving diagnostic accuracy for colorectal conditions in urban settings; however, their adoption and impact on diagnostic outcomes in rural Mozambique's community healthcare settings remain underexplored. A mixed-methods approach combining quantitative data from electronic health records (EHR) with qualitative interviews to understand factors influencing tool adoption and effectiveness. In the study area, digital diagnostics tools were adopted by 45% of community healthcare centers. Diagnostic accuracy improved by an average of 15% when using these tools compared to traditional methods. The use of digital diagnostics significantly enhanced diagnostic outcomes in rural Mozambique's community settings, with a notable improvement in accuracy rates. Further implementation and ongoing evaluation are recommended to sustain the benefits observed and address challenges related to tool accessibility and maintenance. Treatment effect was estimated with  $\text{text}\{logit\}(\pi) = \beta_0 + \beta^{-1} p X_i$ , and uncertainty reported using confidence-interval based inference.

**Keywords:** *Sub-Saharan, Africanization, DigitalHealth, SystemsTheory, Telemedicine, DiagnosticAccuracy, Ethnography*

## ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

**Email:** [info@parj.africa](mailto:info@parj.africa)

Request your copy of the full paper today!

## SUBMIT YOUR RESEARCH

**Are you a researcher in Africa? We welcome your submissions!**

Join our community of African scholars and share your groundbreaking work.

**Submit at:** [app.parj.africa](http://app.parj.africa)



Scan to visit [app.parj.africa](http://app.parj.africa)

**Open Access Scholarship from PARJ**

Empowering African Research | Advancing Global Knowledge