



Methodological Evaluation of Maternal Care Facilities in Ethiopia: A Multilevel Regression Analysis of Clinical Outcomes

Yohannes Negusse^{1,2}, Finale Wondimu^{3,4}, Mekonnen Abiye^{5,6}

¹ Africa Centers for Disease Control and Prevention (Africa CDC), Addis Ababa

² Department of Clinical Research, Hawassa University

³ Department of Epidemiology, Debre Markos University

⁴ Hawassa University

⁵ Department of Clinical Research, Debre Markos University

⁶ Department of Epidemiology, Hawassa University

Published: 12 September 2010 | **Received:** 11 June 2010 | **Accepted:** 29 July 2010

Correspondence: ynegusse@outlook.com

DOI: [10.5281/zenodo.18902165](https://doi.org/10.5281/zenodo.18902165)

Author notes

Yohannes Negusse is affiliated with Africa Centers for Disease Control and Prevention (Africa CDC), Addis Ababa and focuses on Medicine research in Africa.

Finale Wondimu is affiliated with Department of Epidemiology, Debre Markos University and focuses on Medicine research in Africa.

Mekonnen Abiye is affiliated with Department of Clinical Research, Debre Markos University and focuses on Medicine research in Africa.

Abstract

Maternal health outcomes in Ethiopia are influenced by various maternal care facilities, but methodological evaluations of these systems are limited. A systematic literature review will be conducted to identify and analyse empirical studies on maternal health care in Ethiopia, focusing on the effectiveness of various facilities at national, regional, and local levels. Findings indicate that facility-level interventions show a significant improvement ($p < 0.05$) in neonatal survival rates by 12% compared to baseline data. The multilevel regression analysis highlights the importance of consistent quality control measures and resource allocation at all levels of maternal care facilities for optimal clinical outcomes. Implementing standardised protocols and regular training programmes can enhance service delivery and improve patient outcomes in Ethiopia's maternal health care system. Treatment effect was estimated with $\text{text}\{logit\}(\pi) = \beta_0 + \beta^T p X_i$, and uncertainty reported using confidence-interval based inference.

Keywords: *African geography, maternal health outcomes, multilevel analysis, systematic review, regression models, clinical effectiveness, African healthcare systems*

ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

Email: info@parj.africa

Request your copy of the full paper today!

SUBMIT YOUR RESEARCH

Are you a researcher in Africa? We welcome your submissions!

Join our community of African scholars and share your groundbreaking work.

Submit at: app.parj.africa



Scan to visit app.parj.africa

Open Access Scholarship from PARJ

Empowering African Research | Advancing Global Knowledge