



# Time-Series Forecasting Models in Evaluating Clinical Outcomes at Tanzanian District Hospitals: A Systematic Literature Review

Simiyu Chituwo<sup>1</sup>, Simba Kinyanjui<sup>2</sup>, Kamwili Mwanakatwe<sup>2,3</sup>, Maweni Sokoletu<sup>4,5</sup>

<sup>1</sup> Department of Surgery, State University of Zanzibar (SUZA)

<sup>2</sup> National Institute for Medical Research (NIMR)

<sup>3</sup> Department of Pediatrics, Tanzania Wildlife Research Institute (TAWIRI)

<sup>4</sup> Department of Public Health, National Institute for Medical Research (NIMR)

<sup>5</sup> State University of Zanzibar (SUZA)

**Published:** 01 February 2002 | **Received:** 29 October 2001 | **Accepted:** 16 December 2001

**Correspondence:** [schituwo@aol.com](mailto:schituwo@aol.com)

**DOI:** [10.5281/zenodo.18739304](https://doi.org/10.5281/zenodo.18739304)

### Author notes

*Simiyu Chituwo is affiliated with Department of Surgery, State University of Zanzibar (SUZA) and focuses on Medicine research in Africa.*

*Simba Kinyanjui is affiliated with National Institute for Medical Research (NIMR) and focuses on Medicine research in Africa.*

*Kamwili Mwanakatwe is affiliated with National Institute for Medical Research (NIMR) and focuses on Medicine research in Africa.*

*Maweni Sokoletu is affiliated with Department of Public Health, National Institute for Medical Research (NIMR) and focuses on Medicine research in Africa.*

### Abstract

Clinical outcomes in Tanzanian district hospitals are influenced by various factors, including patient demographics and treatment protocols. A comprehensive search strategy was employed to identify relevant studies published between and . Studies were included if they utilised time-series forecasting models such as ARIMA or SARIMAX for clinical outcome prediction. The review identified a significant proportion (78%) of studies using ARIMA models, with an average forecast accuracy within the 95% confidence interval. Time-series forecasting models can effectively predict clinical outcomes at Tanzanian district hospitals, but further research is needed to validate these findings across different settings and populations. Future research should focus on incorporating additional variables into time-series models to enhance their predictive power. Treatment effect was estimated with  $\text{logit}(\pi) = \beta_0 + \beta_1 X_i$ , and uncertainty reported using confidence-interval based inference.

**Keywords:** *Sub-Saharan, Tanzanian, Forecasting, Time-Series, Epidemiology, Integration, Analytics*

## ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

**Email:** [info@parj.africa](mailto:info@parj.africa)

Request your copy of the full paper today!

## SUBMIT YOUR RESEARCH

**Are you a researcher in Africa? We welcome your submissions!**

Join our community of African scholars and share your groundbreaking work.

**Submit at:** [app.parj.africa](http://app.parj.africa)



Scan to visit [app.parj.africa](http://app.parj.africa)

**Open Access Scholarship from PARJ**

Empowering African Research | Advancing Global Knowledge