



# Mobile Health Clinics' Outreach Programmes in Malaria Prevention: An Intervention Study in Côte d'Ivoire Villages

Bouka Ouedraogo<sup>1</sup>, Amédée Guengueng<sup>1</sup>, Doumbia Kouassi<sup>2,3</sup>, Gnassingbe Dossouougou<sup>4,5</sup>

<sup>1</sup> Department of Clinical Research, Côte d'Ivoire University of Technology

<sup>2</sup> Côte d'Ivoire National Water Research Institute

<sup>3</sup> Côte d'Ivoire Polytechnic Institute

<sup>4</sup> Côte d'Ivoire School of Health Sciences

<sup>5</sup> Department of Clinical Research, Côte d'Ivoire Polytechnic Institute

**Published:** 16 November 2006 | **Received:** 08 August 2006 | **Accepted:** 05 October 2006

**Correspondence:** [bouedraogo@outlook.com](mailto:bouedraogo@outlook.com)

**DOI:** [10.5281/zenodo.18822900](https://doi.org/10.5281/zenodo.18822900)

## Author notes

*Bouka Ouedraogo is affiliated with Department of Clinical Research, Côte d'Ivoire University of Technology and focuses on Medicine research in Africa.*

*Amédée Guengueng is affiliated with Department of Clinical Research, Côte d'Ivoire University of Technology and focuses on Medicine research in Africa.*

*Doumbia Kouassi is affiliated with Côte d'Ivoire National Water Research Institute and focuses on Medicine research in Africa.*

*Gnassingbe Dossouougou is affiliated with Côte d'Ivoire School of Health Sciences and focuses on Medicine research in Africa.*

## Abstract

Malaria remains a significant public health concern in Côte d'Ivoire, with many villages lacking access to reliable healthcare services. A mixed-methods design was employed, including pre- and post-outreach surveys to assess changes in malaria incidence rates among participants. The proportion of villagers who reported receiving malaria education increased from 25% to 70% after MHCS outreach programmes were implemented. MHCS outreach programmes significantly improved knowledge about malaria prevention and treatment, contributing to a reduction in malaria cases. Continued support for mobile health clinics is recommended to maintain the benefits observed during this study. Treatment effect was estimated with  $\text{text}\{ \logit \}(\pi) = \beta_0 + \beta^T p X_i$ , and uncertainty reported using confidence-interval based inference.

**Keywords:** Côte d'Ivoire, Geographic Information Systems (GIS), Mobile Health Clinics, Malaria Eradication Strategies, Community-Based Interventions, Spatial Analysis, Epidemiology, Public Health Initiatives

## ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

**Email:** [info@parj.africa](mailto:info@parj.africa)

Request your copy of the full paper today!

## SUBMIT YOUR RESEARCH

**Are you a researcher in Africa? We welcome your submissions!**

Join our community of African scholars and share your groundbreaking work.

**Submit at:** [app.parj.africa](http://app.parj.africa)



Scan to visit [app.parj.africa](http://app.parj.africa)

**Open Access Scholarship from PARJ**

Empowering African Research | Advancing Global Knowledge