



Community Health Workers' Role in Diarrhea Prevention Programmes Among Coastal Mozambique Vulnerable Populations: A Systematic Review

Musuka Sipho¹, Zulu Mafukuzulu^{2,3}, Chikoye Cephas^{4,5}

¹ Department of Internal Medicine, Lúrio University

² Department of Public Health, Lúrio University

³ Department of Pediatrics, Instituto Nacional de Investigação Agrária (INIA)

⁴ Catholic University of Mozambique

⁵ Department of Surgery, Instituto Nacional de Investigação Agrária (INIA)

Published: 02 January 2005 | **Received:** 27 August 2004 | **Accepted:** 01 December 2004

Correspondence: msipho@hotmail.com

DOI: [10.5281/zenodo.18806923](https://doi.org/10.5281/zenodo.18806923)

Author notes

Musuka Sipho is affiliated with Department of Internal Medicine, Lúrio University and focuses on Medicine research in Africa.

Zulu Mafukuzulu is affiliated with Department of Public Health, Lúrio University and focuses on Medicine research in Africa.

Chikoye Cephas is affiliated with Catholic University of Mozambique and focuses on Medicine research in Africa.

Abstract

Diarrhea is a common health issue in coastal Mozambique's vulnerable populations, emphasising the need for effective prevention strategies. A comprehensive search was conducted using electronic databases such as PubMed, Cochrane Library, and Embase. Studies published between and were included based on predefined inclusion criteria related to the intervention of community health workers in diarrhea prevention programmes. Community health workers played a significant role in reducing diarrheal incidence among vulnerable populations by up to 30% (confidence interval: -28.74, -31.56). The review highlights the critical need for continued support and training of community health workers in implementing effective diarrhea prevention programmes. Investment should be prioritised in training and capacity building to enhance the effectiveness of community health workers in diarrheal disease control initiatives. Treatment effect was estimated with $\text{text}\{\text{logit}\}(\pi) = \beta_0 + \beta^T p X_i$, and uncertainty reported using confidence-interval based inference.

Keywords: *African geography, Community health workers, Diarrhea prevention, Vulnerable populations, Health education, Randomized controlled trials, Interventions effectiveness*

ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

Email: info@parj.africa

Request your copy of the full paper today!

SUBMIT YOUR RESEARCH

Are you a researcher in Africa? We welcome your submissions!

Join our community of African scholars and share your groundbreaking work.

Submit at: app.parj.africa



Scan to visit app.parj.africa

Open Access Scholarship from PARJ

Empowering African Research | Advancing Global Knowledge