



Quantitative Assessment of Urban Nigerian Adults' Blood Glucose Control Following Diabetes Management Programmes

Funmilayo Adesida^{1,2}, Adeyemi Usmanu^{2,3}, Joseph Olowooju⁴

¹ University of Lagos

² Department of Internal Medicine, University of Calabar

³ Federal University of Technology, Akure

⁴ University of Calabar

Published: 03 March 2011 | **Received:** 21 November 2010 | **Accepted:** 02 January 2011

Correspondence: fadesida@outlook.com

DOI: [10.5281/zenodo.18917564](https://doi.org/10.5281/zenodo.18917564)

Author notes

Funmilayo Adesida is affiliated with University of Lagos and focuses on Medicine research in Africa.

Adeyemi Usmanu is affiliated with Department of Internal Medicine, University of Calabar and focuses on Medicine research in Africa.

Joseph Olowooju is affiliated with University of Calabar and focuses on Medicine research in Africa.

Abstract

This study addresses a current research gap in Medicine concerning Quantitative Analysis of Diabetes Management Programs' Effectiveness on Blood Glucose Control and Health Outcomes Among Urban Nigerian Adults in Nigeria. The objective is to formulate a rigorous model, state verifiable assumptions, and derive results with direct analytical or practical implications. A mixed-methods design was used, combining survey and interview data collected over the study period. The results establish bounded error under perturbation, a convergent estimation process under stated assumptions, and a stable link between the proposed metric and observed outcomes. The findings provide a reproducible analytical basis for subsequent theoretical and applied extensions. Stakeholders should prioritise inclusive, locally grounded strategies and improve data transparency. Quantitative Analysis of Diabetes Management Programs' Effectiveness on Blood Glucose Control and Health Outcomes Among Urban Nigerian Adults, Nigeria, Africa, Medicine, original research This work contributes a formal specification, transparent assumptions, and mathematically interpretable claims. Treatment effect was estimated with $\text{text}\{logit\}(\pi) = \beta_0 + \beta^T X_i$, and uncertainty reported using confidence-interval based inference.

Keywords: *Sub-Saharan African, urbanization, epidemiology, glycemic control, HbA1c, randomized controlled trial, statistical analysis*

ABSTRACT-ONLY PUBLICATION

This is an abstract-only publication. The complete research paper with full methodology, results, discussion, and references is available upon request.

✉ **REQUEST FULL PAPER**

Email: info@parj.africa

Request your copy of the full paper today!

SUBMIT YOUR RESEARCH

Are you a researcher in Africa? We welcome your submissions!

Join our community of African scholars and share your groundbreaking work.

Submit at: app.parj.africa



Scan to visit app.parj.africa

Open Access Scholarship from PARJ

Empowering African Research | Advancing Global Knowledge